




Dr. Andrew Pippas
 Chief Medical Officer





Angela Dumbuya
 – DNP, NP, AOCNP





Service Area Population Trends






2016: % over age 65*	
Primary Service Area	13.1%
Secondary Service Area	12.0%
Total	12.6%







Who is a cancer survivor

- The IOM in 2006 defines survivorship as the "phase of care that follows primary treatment
- Memorial Sloan Kettering defines survivorship as the transition from cancer patient to cancer survivor this occurs after you have completed active treatment, as determined by your treating physician.
- According to MD Anderson a survivor is anyone who's been diagnosed with cancer. which starts at the time of diagnosis and continues throughout the rest of the patient's life.
- The NCI defines a survivor as one diagnosed with a malignancy and this
 individual would be considered a survivor from the time of diagnosis through
 the balance of his or her life.

 IOM Recommended Components of Survivorship Care

PreventionOf recurrent

and new cancers and late effects

Surveillance

For metastasis, recurrence, or 2nd cancers; assessment of medical & psychosocial late effects

impacts of cancer and its treatment

Coordination

between specialists and primary care providers

 Hewitt M, Greenfield S, Stovall E. From Cancer Patient to Cancer Survivor: Lost in Transition. 2005. Washington, D.C.: The National Academies Press.

- The Accreditation Committee made the following changes to the established time frame and scope of implementation for Standard 3.3.
- January 1, 2015 Implement a pilot survivorship care plan process involving 10% of eligible patients.
 - January 1, 2016 Provide survivorship care plans to 25% of eligible patients.
 - January 1, 2017 Provide survivorship care plans to 50% of eligible patients.
 - January 1, 2018 Provide survivorship care plans to 75% of eligible patients.
 - January 1, 2019 Provide survivorship care plans to all eligible patients.

Cancer Programs that have fully implemented the Standard by the time of their on-site visit during the 2015, 2016, 2017 survey cycle, will receive special recognition in their Performance Reports at the time of their next survey.

- Survivorship Program at John B. Amos Cancer Center. (JBACC)
 - Established in September of 2014.
 - It is a consultative/shared-care model
 - Program headed by Chief Medical Director Dr. Pippas.
 - Program run by NP Angela Dumbuya DNP.
 - Patients are referred to the NP after adjuvant/initial therapy (excluding hormonal therapy) for short term follow up. Patients are also seen in the program after five years of follow up with oncologist.

Who is a cancer survivor at JBACC?

- An adult patient diagnosed with a malignancy and has completed active treatment as determined by the treating physician and has no evidence of disease. Survivorship continues for the rest of the patient's life.
- All patients, meeting the above criteria are referred to the survivorship program six weeks after completion of primary therapy (Hormone therapy not included).
- Referrals to the program can also be done after surgery for patients not needing chemotherapy or XTR. Patients can self refer from outside Center who have been treated by other oncologists within our region.

Referral/Consultative Model

- Patients are referred to the survivorship program for a one time visit to meet with the NP
 - Patient undergoes a history and physical exam
 - Evaluate and manage patient's sequelae of treatment (medical, nutritional and psychosocial).
 - Educate patient about survivorship issues.
 - Discuss and refer for other screening tests (colonoscopy, mammogram, PSA, lung cancer screening, skin/dermatology).
 - Make referral to health promotion (smoking cessation, nutrition, sexual, social and psychological health, rehabilitation and exercise).
 - Care plan and treatment summary are reviewed with patient.
 - Send patient back to treating oncologist to continue follow up care at the end of the visit.
 - Permanent transfer to NP run survivor program is at the discretion of the treating oncologist. Most of our PCPs do not want to see patients for cancer follow-up.
 - Follow NCCN guidelines.

The CARE Plan Report

Survivorship Care Plan

- Since the publication of <u>From Cancer Patient to Cancer Survivor: Lost in</u> Transition care plans have received lots of attention
- It is a record of a patient's cancer treatment and recommendations for followup care.
- IOM strongly recommends that at the completion of cancer treatment, clinicians provide patients and primary care providers with a summary of the treatment delivered and a detailed plan for ongoing care, including follow-up schedules for visits and testing, and recommendations for the early detection and management of treatment-related effects and other health problems.
- Elements of a Care Plan (Treatment summary, follow up care and education material regarding potential late effects of cancer therapy).
- JBACC: Equicare is the choice of our survivorship care plan. This generates information from our MHR, Aria. Difficult to separate demographics.

Do Treatment Summaries Work?

- The Oncologist, 2016: UAB/CCN, (Kenzik et al): Treatment Summaries and Follow-Up Care Instruction for Cancer Survivors: Improving Survivor Self-Efficacy and Health Care Utilization.
- 441 survivors, >65 years old, 12 centers
- "Self-Efficacy": Stanford Chronic Illness scale measuring a person's confidence in his/her ability to mange health conditions caused by cancer/treatment.
- Contacted at home
 - 40% written summary
 - 35% follow up care plan
 - 79% had a verbal explanation of care plan
- All associated with high 'self-efficacy' scores: patients able to manage their chronic illnesses better and had fewer ER visits.

- Survivorship Program at JBACC
 - 401 patient have been seen in the survivorship program since September of 2014.
 - 327 have been breast cancer patients.
 - 44 colorectal cancer
 - 10 head and neck cancer
 - 10 lymphoma
 - 3 Gastric
 - 7 others.

- Survivorship Program at JBACC
 - 2016 Update:
 - 117 patients have been seen.
 - 92 breast cancer patients.
 - 13 colorectal patients
 - 6 Lymphoma
 - 2 Gastric
 - 1 Lung
 - I Liver
 - 1 Multiple Myeloma
 - 2 Head & Neck.
 - One patient had 2 separate diagnoses of breast and colon.

National Survivorship Spectrum: Similar to JBACC

Statistics on survivors




Figure 1: Estimated number of cancer survivors in the United States by disease site as of January 1, 2005 (N = 11.1 million). Data from Reference 3. Retrieved from: http://dccps.nci.nih.gov/ocs/prevalence/prevalence.html#survivor

Issues facing survivors

- Physical
- Insomnia
- Emotional
- Practical Employment
- Osteoporosis
- Lymphedema
- Menopausal symptoms
- Neuropathy
- Second malignancy
- Fertility/sexuality
- Cognitive
- Hair/skin/nail
- Pain

- Barriers facing survivors.
 - Fragmented healthcare delivery: may need to see multiple providers.
 - Cost: All of our patients complain of facility fee costs.
 - Barriers to access: getting to our center may be a challenge. Serve 14 counties in our primary market. 40% of patients drive ~30 miles to get to JBACC.
- Barriers facing providers.
 - Fragmented system of care: Should this patient go back to PCP? Is this
 patient best served by seeing PCP or having serial evaluation by oncology?
 Difference between a testicular cancer patient and Stage 1 breast cancer
 patient.
 - System does NOT automatically refer to Survivorship.
 - Evolving survivorship standards of care: what about addressing metabolic/weight issues: who makes referral to bariatric clinic?
 - Difficulties in roles between providers. PCPs vary in their expectations of surveillance. Manage "just the cancer" or general health?

Services Provided

- Psychosocial Services; initial intake evaluation for every patient. In survivorship, seen on an individual basis(by D. Johnson)
- Nutritional services (Beth Bussey)
- Rehabilitation (PT/OT/lymphedema)
- Sexual Health (D. Johnson)
- Genetic counseling (Cindy Snyder, RN. St Francis)
- Fertility (before and after chemo/XRT. Dr. Thiruppathi)
- Patient Portal (labs, office visits)
- Support groups
 - Man to Man, Woman to Woman, Next Step, Look Good Feel Better, T'ai Chi
- Skin care (late effects of paronychia, radiation mastitis)
- Chronic pain management (referral to specialty practice)
- Exercise: group program
- Smoking cessation
 - Group (D Johnson, M Knobo)
 - If a pharmacological intervention is needed then patient sees the NP.

Smoking Cessation: Onsite facilitator

- Freedom From Smoking Group=learn coping skills to stop smoking. Reported 29% I year quit rate in a 1996 overview of success rates.
- Nicotine more addictive than opiates, alcohol
- 1st cycle 8/17/16, next cycle 11/2/16
 - 46 referrals
 - 6 participated in the 7 week program
 - 1 'graduated' and stopped smoking—for now...

"The Spirit is willing, the Flesh is trying"

Recent New Programs

- Art Therapy: meet monthly to work of a collage for JBACC.
- Massage: "Massage is good medicine."
 Massage therapist; may be useful reducing treatment/surgical pain and cancer pain. May improve immune system.
- Pet Therapy: part of hospital program.

Thoracic Oncology: LDSCT

- Dr. Peter Seirafi: 11/20/13. Lung cancer navigator/team.
- Screened high risk smokers (30 pk/years, 55yo)
- 252 patients, 12 asymptomatic lung cancer patients.
 - 4.79% similar to the LDSCT/ACRIN trial of 3.97 %

Nutritional Health

- Evaluate all patients
- LAUNCH program for breast survivors: to be replaced with the LIFT (live inspired, fight together) program through WW.
- Manage long term TPN, enteral nutrition
- Oversee/discuss complementary therapy
 - 80% of patients use alternative treatments
- Focus on Head and Neck/Thoracic/Esophageal and GI surgical patients

- Additional Services.
 - Long term educational workshop that may be provided for our patients.
 - E Learning
 - Nutrition: Eating well after cancer
 - PTSD in cancer patients
 - Psychosocial distress in cancer patients
 - Physician and patients' perspective in cancer treatment
 - Neuropathy care for cancer survivors
 - Exercise for survivors
 - Long term complication after XRT
 - Lymphedema
 - Improving cognitive functions after treatment
 - Fertility and sexual health after cancer treatment
 - Long-term effects of transplantation


Survivorship Research Studies

- EROS: Engendering Reproductive Health within Oncologic Survivorship.
 - Females 18-55 years, with intact fertility, assessing reproductive and sexual health
- Alliance 22102: Topical testosterone/placebo in the adjuvant treatment of AI arthralgias.
- SWOG 0820: Preventing Adenomas of the Colon with Eflornithine and Sulindac in ST 0-3 colon cancer.
- Alliance 11401: Role of weight loss in adjuvant treatment of overweight(BMI </=27)/obese women with breast cancer.
 - 2 year Health education +/-supervised weight loss(Fitbit, phone instruction BWEL center DFCI)


OBESITY: The GREAT BATTLE

- "We kill the cows, and the cows kill us."
- Breast cancers survivors are twice as likely to gain weight and non cancer patient(ACS)
 - During treatment, 80% of woman gain weight.


Weight and Survival in Early-Stage Breast Cancer



Nurses' Health Study: Physical activity after breast cancer diagnosis




The HEAL Study: Physical Activity after Diagnosis




Irwin ML et al JCO, 2008.

Summary of Mortality from Cancer According to Body-Mass Index for U.S. Men in the Cancer Prevention Study II, 1982 through 1998





Mortality from Cancer According to BMI for U.S. Women in the Cancer Prevention Study II



Cancer Site or Type
Strength of the Evidence
in Humans†
Relative Risk of the Highest
BMI Category Evaluated
versus Normal BMI (95% CI)‡

Esophagus: adenocarcinoma Sufficient 4.8 (3.0–7.7)

Gastric cardia Sufficient 1.8 (1.3–2.5)

Colon and rectum Sufficient 1.3 (1.3–1.4)

Liver Sufficient 1.8 (1.6–2.1)

Gallbladder Sufficient 1.3 (1.2–1.4)

Pancreas Sufficient 1.5 (1.2–1.8)

Breast: postmenopausal Sufficient 1.1 (1.1–1.2)§

Corpus uteri Sufficient 7.1 (6.3-8.1)

Ovary Sufficient 1.1 (1.1–1.2)

Kidney: renal-cell Sufficient 1.8 (1.7–1.9)

Meningioma Sufficient 1.5 (1.3–1.8)

Thyroid Sufficient 1.1 (1.0-1.1)§


Multiple myeloma Sufficient 1.5 (1.2–2.0)

NYT: August 2016

- Strong evidence was already available to link five cancers to being overweight or obese: adenocarcinoma of the esophagus; colorectal cancer; breast cancer in postmenopausal women; and uterine and kidney cancers.
- This new review, published in The New England Journal of Medicine, links an
 additional eight cancers to excess fat: gastric cardia, a cancer of the part of the
 stomach closest to the esophagus; liver cancer; gallbladder cancer; pancreatic
 cancer; thyroid cancer; ovarian cancer; meningioma, a usually benign type of brain
 tumor; and multiple myeloma, a blood cancer.
- According to the chairman of the working group, Dr. Graham Colditz, a professor of medicine and surgery at Washington University in St. Louis, these 13 cancers together account for 42 percent of all new cancer diagnoses.
- "Only smoking comes close" as an environmental factor affecting cancer risk, Dr.
 Colditz said. "And that's an important message for nonsmokers. Obesity now goes
 to the top of the list of things to focus on."

Benefits of Increasing Post-Diagnosis PA

- Need to encourage cancer survivors to maintain (if active prior to diagnosis) or increase physical activity after diagnosis.
- Physical activity is associated with numerous benefits
 - Decreasing adverse side effects of surgery and treatment
 - Improving quality of life and fatigue
 - Improving fitness, functional capacity, and body composition
 - Decreasing serum hormones and growth factors
 - Improving cancer-specific survival and all cause survival



Telegraph.co.uk

Home News Sport Finance Lifestyle Comment Travel Culture Technology **Politics** World Celebrities Obituaries Weird Earth Science Health News Education Royal Family Road and Rail Defence Scotland Northern Ireland Religion

HOME - NEWS - UK NEWS

Soaring obesity 'will cause cancer time bomb'

By Stephen Adams Published: 12:01AM BST 21 May 2007



40 per cent of cancers can be prevented through diet and exercise

Urgent action is needed to prevent a "cancer time bomb" exploding as a result of soaring obesity rates, a leading specialist warns today.



Fund.